


276
277
278
279
280
281
282
283
284
286
287
288
289
290
291
292
293
294
295
296
297
298

R. J. Klaunig Ruch. Inhibition of mouse hepatocyte intercellular communication by paraquat-generated oxygen free radicals. <i>Toxicol Appl Pharmacol.</i> 1988. 94:427-36
J. Sela Starr. Resistance to paraquat in a mammalian cell line. <i>Somat Cell Mol Genet.</i> 1986. 12:141-52
D. Sandy Di Monte. Comparative studies on the mechanisms of paraquat and 1-methyl-4-phenylpyridine (MPP+) cytotoxicity. <i>Biochem Biophys Res Commun.</i> 1986. 137:303-9
A. F. Jornot Junod. Comparative study on the selenium- and N-acetylcysteine-related effects on the toxic action of hyperoxia, paraquat and the enzyme reaction hypoxanthine-xanthine oxidase in cultured endothelial cells. <i>Agents Actions.</i> 1987. 22:176-83
T. M. Brown Hagen. Protection against paraquat-induced injury by exogenous GSH in pulmonary alveolar type II cells. <i>Biochem Pharmacol.</i> 1986. 35:4537-42
C. Junod Ody. Direct toxic effects of paraquat and oxygen on cultured endothelial cells. <i>Lab Invest.</i> 1985. 52:77-84
J. J. Kimura Huang. A specific role of reduced adrenodoxin in adrenal mitochondrial steroid hydroxylases. <i>Biochem Biophys Res Commun.</i> 1971. 44:1065-70
J. Wei Sinow. Ocular toxicity of paraquat. <i>Bull Environ Contam Toxicol.</i> 1973. 9:163-8
M. Joyce. Ocular damage caused by Paraquat. <i>Br J Ophthalmol.</i> 1969. 53:688-90
I. Nemcsok Rojik. Morphological and biochemical studies on liver, kidney and gill of fishes affected by pesticides. <i>Acta Biol Hung.</i> 1983. 34:81-92
J. S. Gibson Bus. Paraquat: model for oxidant-initiated toxicity. <i>Environ Health Perspect.</i> 1984. 55:37-46
Y. Tabor Shiloh. Abnormal response of ataxia-telangiectasia cells to agents that break the deoxyribose moiety of DNA via a targeted free radical mechanism. <i>Carcinogenesis.</i> 1983. 4:1317-22
J. R. Norbeck Dawson. The effectiveness of N-acetylcysteine in isolated hepatocytes, against the toxicity of paracetamol, acrolein, and paraquat. <i>Arch Toxicol.</i> 1984. 55:11-5
E. Brigelius Cadenas. Paraquat-induced chemiluminescence of microsomal fractions. <i>Biochem Pharmacol.</i> 1983. 32:147-50
R. J. Swoboda Brennan. Oxidative mutagens induce intrachromosomal recombination in yeast. <i>Mutat Res.</i> 1994. 308:159-67
P. Petronilli Costantini. On the effects of paraquat on isolated mitochondria. Evidence that paraquat causes opening of the cyclosporin A-sensitive permeability transition pore synergistically with nitric oxide. <i>Toxicology.</i> 1995. 99:77-88
S. Ona Kojima. Antioxidative activity of 5,6,7,8-tetrahydrobiopterin and its inhibitory effect on paraquat-induced cell toxicity in cultured rat hepatocytes. <i>Free Radic Res.</i> 1995. 23:419-30
T. H. Chu Tseng. Crocetin protects against oxidative damage in rat primary hepatocytes. <i>Cancer Lett.</i> 1995. 97:61-7
M. L. Lee Kuo. Pronounced activation of protein kinase C, ornithine decarboxylase and c-jun proto-oncogene by paraquat-generated active oxygen species in WI-38 human lung cells. <i>Biochim Biophys Acta.</i> 1995. 1268:229-36
M. L. Lin Kuo. The genotoxicity of the waste water discharged from paraquat manufacturing and its pyridyl components. <i>Mutat Res.</i> 1993. 300:223-9
M. Eguchi-Kasai Murakami. Biological effects of active oxygen on an X-ray-sensitive mutant mouse cell line (SL3-147). <i>Mutat Res.</i> 1995. 336:215-21
U. Glatzle Koch. Measurement of ion leakage from plant cells in response to aquatic pollutants. <i>Bull Environ Contam Toxicol.</i> 1995. 54:606-13

Intercellular communication through gap junctions functions in electrical synapsing, homeostasis, hormonal response, en
Paraquat-resistant variants were isolated in Chinese hamster ovary (CHO) cells by stepwise increases in paraquat concen
1-methyl-4-phenylpyridine (MPP+) is the putative toxic metabolite of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MP
The potential protective effect of N-acetylcysteine against various types of oxidative stress (exposure to hyperoxia, treat
Exogenous GSH provided rat alveolar type II cells with significant protection against injury induced by paraquat. This prot
Paraquat (PQ) is a herbicide known to generate O2 radicals and to injure lung epithelial cells, leading eventually to pulmo
Effect of a herbicide, paraquat (1,1'-dimethyl-4,4'-bipyridilium-dichloride), the fungicide copper sulphate, and zinc chlorid
Paraquat, a quaternary ammonium bipyridyl herbicide, produces degenerative lesions in the lung after systemic adminis
A defect in DNA repair coupled to anomalous DNA synthesis after induction of certain radiogenic DNA damage is suspect
The protective effect of N-acetylcysteine against the toxicity of paracetamol, acrolein, and paraquat was investigated usi
Active oxygen species are thought to be involved in the causation of a number of diseases including cancers. We have inv
This paper reports an investigation on the effects of the bipyridylum herbicide, paraquat, on rat liver mitochondria in vit
The in vitro antioxidative activity of 5,6,7,8-tetrahydrobiopterin (BPH4) was measured and the ability of BPH4 to prevent
Crocetin is a major component in the fruit of Gardenia jaminoides Ellis, a Chinese herbal medicine. Its protective action a
Paraquat (methyl viologen, PQ) is a widely used herbicide that produces oxygen-derived free radicals and severely injure
The genotoxic potencies of waste water condensate from paraquat manufacturing and its pyridyl components on V79 cel
The biological effects of active oxygen species were examined in a mutant mouse cell line (SL3-147) that is deficient in th

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant





300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
324

L. J. Craig Walker. A role for the human DNA repair enzyme HAP1 in cellular protection against DNA damaging agents and hypoxic stress. <i>Nucleic Acids Res.</i> 1994. 22:4884-9
T. Reinheckel Grune. Proteolysis in cultured liver epithelial cells during oxidative stress. Role of the multicatalytic proteinase complex, proteasome. <i>J Biol Chem.</i> 1995. 270:2344-51
D. Amirov Ostrovsky. Bacteria and pesticides: a new aspect of interaction--involvement of a new biofactor. <i>Biofactors.</i> 1994. 4:151-4
N. Bowles Chen. Prevention of paraquat toxicity in suspensions of alveolar type II cells by paraquat-specific antibodies. <i>Hum Exp Toxicol.</i> 1994. 13:551-7
C. M. Moreno Palmeira. Metabolic alterations in hepatocytes promoted by the herbicides paraquat, dinoseb and 2,4-D. <i>Arch Toxicol.</i> 1994. 68:24-31
T. C. Lai Lee. Protection of a rat tracheal epithelial cell line from paraquat toxicity by inhibition of glucose-6-phosphate dehydrogenase. <i>Biochem Pharmacol.</i> 1993. 45:1143-7
H. Palace Babich. Oxidative stress in fish cells: in vitro studies. <i>Arch Environ Contam Toxicol.</i> 1993. 24:173-8
B. J. Shawen Day. A metalloporphyrin superoxide dismutase mimetic protects against paraquat-induced endothelial cell injury, in vitro. <i>J Pharmacol Exp Ther.</i> 1995. 275:1227-32
C. M. Moreno Palmeira. Thiols metabolism is altered by the herbicides paraquat, dinoseb and 2,4-D: a study in isolated hepatocytes. <i>Toxicol Lett.</i> 1995. 81:115-23
B. D. Joshi Uhal. Fibroblasts isolated after fibrotic lung injury induce apoptosis of alveolar epithelial cells in vitro. <i>Am J Physiol.</i> 1995. 269:L819-28
E. Baum Hoffer. N-acetylcysteine increases the glutathione content and protects rat alveolar type II cells against paraquat-induced cytotoxicity. <i>Toxicol Lett.</i> 1996. 84:7-12
M. Gross Poot. Cell cycle defect in connection with oxygen and iron sensitivity in Fanconi anemia lymphoblastoid cells. <i>Exp Cell Res.</i> 1996. 222:262-8
L. Kelsey Jia. Increased activity and sensitivity of mitochondrial respiratory enzymes to tumor necrosis factor alpha-mediated inhibition is associated with increased cytotoxicity in drug-resistant leukemic cell lines. <i>Blood.</i> 1996. 87:2401-10
T. Reinheckel Grune. Degradation of oxidized proteins in K562 human hematopoietic cells by proteasome. <i>J Biol Chem.</i> 1996. 271:15504-9
G. B. Gwinner Stephanz. Heat-aggregated IgG and interleukin-1-beta stimulate manganese superoxide dismutase in mesangial cells. <i>Exp Nephrol.</i> 1996. 4:151-8
T. Imaeda Tsuchiya. Detoxification of paraquat poisoning: effects of carbohydrate sulfate, alkylsulfate and alkylsulfonate on active oxygen. <i>Biol Pharm Bull.</i> 1995. 18:1700-4
M. Day Patel. Requirement for superoxide in excitotoxic cell death. <i>Neuron.</i> 1996. 16:345-55
S. Jain Ali. Paraquat induced DNA damage by reactive oxygen species. <i>Biochem Mol Biol Int.</i> 1996. 39:63-7
C. D. Rice Lindsay. Assessment of the biochemical effects of percutaneous exposure of sulphur mustard in an in vitro human skin system. <i>Hum Exp Toxicol.</i> 1996. 15:237-44
J. Yamanaka Kitahara. Mutagenicity of cobalt and reactive oxygen producers. <i>Mutat Res.</i> 1996. 370:133-40
S. H. Tsai Jee. Bipyridyl dihydrochloride inhibits tumor necrosis factor-alpha secretion by human keratinocytes on ultraviolet irradiation. <i>J Formos Med Assoc.</i> 1996. 95:706-8
G. Maggioni Cappelletti. Dynamic changes in the state of actin polymerization in human alveolar cells exposed to the oxidant agent paraquat. <i>Eur J Cell Biol.</i> 1996. 71:293-302
H. Siow Sato. Expression of stress proteins heme oxygenase-1 and -2 in acute pancreatitis and pancreatic islet betaTC3 and acinar AR42J cells. <i>FEBS Lett.</i> 1997. 405:219-23

The HAP1 protein (also known as APE/Ref-1) is a bifunctional human nuclear enzyme required for repair of apurinic/apyr
Exposure to various forms of mild oxidative stress significantly increased the intracellular degradation of both "short-live
Positively charged hydrophobic pesticides of the dipyridyl family [diquat, paraquat, benzylviologen (BV++), etc.] were sho
1. The herbicide, paraquat, is accumulated by the energy-dependent polyamine uptake pathway of alveolar type II cells.
The cytotoxic effects of the herbicides paraquat (1,1'-dimethyl-4,4'-bipyridylium dichloride), dinoseb (2-sec-butyl-4,6-din
Transformed rat tracheal epithelial cells (U2) were found to be 3.6-fold more sensitive than lung fibroblasts (RLF) to para
Bluegill sunfish BF-2 fibroblasts were used in the neutral red (NR) cytotoxicity assay to discern the toxicities of hydrogen
There is an increased interest in development of therapeutic agents to treat disease states that involve reactive oxygen s
This report is an extension and complement of a previous study reporting the effect of three herbicides (paraquat, dinose
Primary lung fibroblasts were isolated from patients with idiopathic pulmonary fibrosis (HIPF), from normal human lung
A protective effect of N-acetylcysteine in oxidative lung damage was reported by a number of workers; however, the me
Fanconi anemia (FA) is an autosomal recessive disorder involving progressive pancytopenia, skeletal malformations, and
The drug-resistant leukemic cell lines, CEM/VLB100 and K/DAU600, are more sensitive to tumor necrosis factor alpha (TN
Exposure to various forms of oxidative stress (H2O2 and O2.-) significantly increased the intracellular degradation of both
The response of the glomerular mesangial cell to reactive oxygen species (ROS) has been implicated in the pathogenesis
The potency of carbohydrate sulfate (dextran sulfate, DS; glucose sulfate, GS), alkyldisulfonate (butane-disulfonate, BDS;
We tested the pathogenic role of O2-) radicals in excitotoxic injury. Inactivation of the TCA cycle enzyme, aconitase, was
The redox cycling contact herbicide paraquat (PQ) causes oxidative damage to pulmonary tissue. PQ is reduced enzymati
1. Sulphur mustard (HD) is a potent chemical warfare agent which causes incapacitating blisters on human skin. There is
Oxidative stress has been implicated in carcinogenesis yet there are chemicals that produce oxidative stress that are not
Paraquat manufacturers in Taiwan have been found to develop solar lentigo, actinic keratosis, as well as skin cancer in su
To investigate the role of dynamic changes in actin cytoskeleton in cellular response to oxidative stress, we have analyze
Oxygen-derived free radicals have been implicated in the pathogenesis of acute pancreatitis, yet adaptive responses in th

